




Neuromyelitis Optica Spectrum Disorder in Active Duty Service Members



Jon P. Williams, DO, PhD^{1,2,3}, Jonathan Galli, MD^{1,2}, Keith A. Groshans MD⁴, David E. Horvat MD⁴, Sharon Austin PhD², Sean J. Pittcock MD⁵, John W. Rose MD^{1,6}, Noel G. Carlson PhD^{1,6,7}, John E. Greenlee MD^{1,2}, M. Mateo Paz Soldan MD PhD^{1,2}, Nathan T. Tagg^{4,8}, and Stacey L. Clardy MD PhD^{1,2,6}

1. University of Utah, Department of Neurology, Salt Lake City, UT. 2. George E. Wahlen Department of Veterans Affairs Medical Center, Salt Lake City, UT. 3. US Air Force Institute of Technology, 4. Walter Reed National Military Medical Center 5. Mayo Clinic Neuroimmunology Laboratory, Rochester, MN, 6. University of Utah, School of Medicine, Salt Lake City, UT. 7. University of Utah, Department of Neurobiology and Anatomy, Salt Lake City, UT. 8. Uniformed Services University of Health Sciences



SALT LAKE CITY

Objective

To characterize patients with neuromyelitis optica spectrum disorder (NMOSD) in the Department of Defense (DoD) population

Background

- NMOSD is an astrocytopathy with an associated serum autoantibody to the aquaporin 4 water channel resulting in secondary demyelination^{1,2}
- NMOSD preferentially involves the optic nerves, spinal cord and circumventricular organs
- NMOSD is associated with additional comorbid autoimmunity^{3,4}
- The incidence of NMOSD among military personnel is unknown; the overlap of NMOSD with other autoimmune conditions has not been defined in this population

Design and Methods

- Approved U of Utah/VA IRB # 90978
- Comprehensive query of patient records in the DoD for relevant diagnostic codes (ICD-9 341.0 and ICD-10 G36.0) between Jan 1 2010 and August 1, 2017
- Review of patient records via Joint Legacy Viewer, AHLTA, and Essentris, the DoD electronic medical record

Funding

The authors are grateful for the support of the Transverse Myelitis Association (TMA) and Barbara Gural-Steinmetz

Results

- Identified **131** unique patients within the DoD system with a ICD code documented in the medical record at least once
- Of these patients, 39 were confirmed as service members; the remainder were dependent beneficiaries
 - 17 met 2015 diagnostic criteria for NMOSD⁵
 - Patients were categorized as:
 - NMOSD with documented AQP-IgG seropositivity: 15
 - Seronegative NMOSD: 2
 - 22 did not meet diagnostic criteria for NMOSD:
 - Alternative diagnoses: 15
 - Insufficient evidence to determine diagnosis: 7
 - Of the **17** patients meeting diagnostic criteria for NMOSD:
 - Sex:**
 - Female: 9
 - Male: 8
 - Ethnicity:**
 - African American: 6
 - Caucasian: 6
 - Asian: 1
 - Hispanic/ Latino: 1
 - Other: 2
 - Unknown: 1
 - Average age of onset:** 39.3
 - Range of ranks from junior enlisted (E3) to officers (O4)
 - DoD follow up ranged from 6 to 168 months
 - Average modified Rankin scale (mRS) at final follow up: 2
 - Of the 22 patients not meeting diagnostic criteria diagnoses included:
 - Demyelinating disease/ multiple sclerosis (7), isolated myelitis (4), isolated optic neuritis (3), mixed connective tissue disorder (1), insufficient data (7)

Figure 1: Clinical Events




Figure 1. Clinical events in DoD patients diagnosed with NMOSD. Top panel is presenting symptoms. Middle and bottom panels represent distribution of longitudinally extensive transverse myelitis, and documented cases of optic neuritis, respectively, at any time since presentation. The x-axis is number of patients for all panels.

Figure 2: CSF characteristics




Figure 2. Cerebrospinal fluid (CSF) analysis of DoD patients meeting diagnostic criteria for NMOSD. OCB, oligoclonal bands.

Figure 4: Clinical outcomes




Figure 4. Modified Rankin Scale at most recent follow up for DoD patients meeting diagnostic criteria for NMOSD.

Figure 3: Treatment exposure




Figure 3. Treatment exposure of DoD patients meeting diagnostic criteria for NMOSD. PLEX, plasma exchange; IVIG, intravenous immunoglobulin.

Conclusions

- This is a first-ever characterization NMOSD in the DoD population.
- Demographics and clinical characteristics agree with prior reports⁶, with the exception of sex ratio, which may reflect a unique influence of this population.
- The study is limited by the time range (2010–2017) of analysis, and will benefit from further evaluation of additional DoD cases prior to 2010.

References

1. Lennon VA et al. A serum autoantibody marker of neuromyelitis optica: distinction from multiple sclerosis. *Lancet*. 2004 Dec 11-17;364(9451):2106-12.
2. Papadopoulos MC, Verkman AS. Aquaporin 4 and neuromyelitis optica. *Lancet Neurol*. 2012 Jun;11(6):535-44.
3. Leite MI et al. Myasthenia gravis and neuromyelitis optica spectrum disorder: a multicenter study of 16 patients. *Neurology*. 2012 May 15;78(20):1601-7.
4. Wingerchuk DM, Weinshenker BG. The emerging relationship between neuromyelitis optica and systemic rheumatologic autoimmune disease. *Mult Scler*. 2012 Jan;18(1):5-10.
5. Wingerchuk, DM et al. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders *Neurology*. 2015 Jul 14; 85(2): 177–189.
6. Kim SH et al. Clinical spectrum of CNS aquaporin-4 autoimmunity. *Neurology*. 2012 Apr;78(15):1179-85.